

Gravitational Microlensing Influence on X-ray Radiation from Accretion Disk of Active Galaxies

Predrag Jovanović

pjovanovic@aob.bg.ac.yu

Astronomical Observatory, Volgina 7, 11160 Belgrade, Serbia and Montenegro

ABSTRACT

Gravitational microlensing is a very useful tool for investigating the innermost part of lensed quasars, especially for studying a relativistic accretion disk around a massive black hole (BH) supposed to exist in a quasar's center. Here we present a short overview of our recent investigations in this field.

Subject headings: accretion, accretion disks — line: profiles — galaxies: active — gravitational lensing — X-rays: galaxies

The aim of the dissertation was to investigate the influence of gravitational microlensing on detected X-ray radiation from accretion disks of active galactic nuclei (AGNs). For this purpose, the focus was set to the analysis of Fe $K\alpha$ spectral line and X-ray continuum variations due to gravitational microlensing. Besides, a significant part of the dissertation was devoted to the optical depth of microlensing (i.e. to the probability to observe these effects) and to the corresponding time scales.

The initial assumption was the existence of a super massive black hole ($10^7 - 10^9 M_\odot$), surrounded by an accretion disk that radiates in X-rays, in the center of all types of AGNs. Accretion disks could have different forms, dimensions and emission, depending on the type of central BH, being rotating (Kerr metric) or non-rotating (Schwarzschild metric). Except for effects due to disk instability, its emission could also be affected by gravitational microlensing, especially in the case of the gravitationally lensed quasars (Chartas et al. 2002; Dai et al. 2003; Chartas et al. 2004).

The disk emission was analyzed by numerical simulations, based on a ray-tracing method in a Kerr metric, taking into account only photon trajectories reaching the observer's sky plane (see Popović et al. (2003b) and references therein). The influence of microlensing on a standard accretion disk was studied using three types of a microlensing model: point-like microlens, straight-fold caustic and quadruple microlens (microlensing pattern). An

illustration of the straight-fold caustic crossing over an accretion disk in the Kerr metric and the corresponding effects on the shapes of the X-ray continuum and the Fe $K\alpha$ line are shown in Fig. 1.

Some conclusions of the dissertation are:

1. Gravitational microlensing can produce significant variations and amplifications of the line and continuum fluxes. These deformations of the X-ray radiation depend on both the disk and microlens parameters. For more details see Popović et al. (2001, 2003a)
2. Even very small mass objects could produce noticeable changes in the Fe $K\alpha$ line profile during a microlensing event. This is because of the small dimensions of the X-ray emitting region (Popović et al. 2003b; Popović & Jovanović 2002)
3. Microlensing can satisfactorily explain the excess in the iron line emission observed in three gravitational lens systems: MG J0414+0534 (Chartas et al. 2002), QSO 2237+0305 (Dai et al. 2003) and H 1413+117 (Chartas et al. 2004). More details can be found in Jovanović et al. (2003); Jovanović (2005); Jovanović & Popović (2005) and Popović et al. (2003b, 2006)
4. On the basis of these investigations, one can expect that the Fe $K\alpha$ line and X-ray continuum amplification due to microlensing can be significantly larger than the corresponding effects on optical and UV emission lines and continua (see Fig. 2 and Popović et al. (2002a,b, 2003b); Jovanović et al. (2006))
5. The optical depth for microlensing of X-ray emitting region by compact objects from the bulge/halo of the host galaxy is very small, $\tau \sim 10^{-4}$ (Popović et al. 2002a, 2003b; Zakharov et al. 2004)
6. The optical depth for gravitational microlensing by cosmologically distributed deflectors could be significant and could reach $\tau \sim 0.01 - 0.1$. In that case, the maximum optical depth ($\tau \sim 0.1$) could be expected if dark matter forms cosmologically distributed compact objects (see Zakharov et al. (2004, 2005))
7. Although microlensing is an achromatic effect, it can induce wavelength dependent variations of the X-ray continuum of $\sim 30\%$ due to the radial distribution of temperature in an accretion disk (Popović et al. 2006)
8. Analysis of three different microlensing time scales (see Fig. 2) led to the conclusion that the variations of the X-ray continuum are the fastest ($\sim$ several months) in comparison to the variations of the optical and UV continua, which are weaker and much slower ($\sim$ several years) (Jovanović et al. 2006)

According to the results of the dissertation, one can conclude that monitoring of gravitational lenses in the X-ray spectral range may help us to understand the physics and to reveal the structure of the innermost parts of active galaxies, particularly their relativistic accretion disks.

This work is a part of the project: "Astrophysical Spectroscopy of Extragalactic Objects" supported by the Ministry of Science, Technologies and Development of Serbia.

REFERENCES

- Chartas, G., Agol, E., Eracleous, M., Garmire, G., Bautz, M. W., Morgan, N. D. 2002, ApJ, 568, 509
- Chartas, G., Eracleous, M., Agol, E., Gallagher, S. C. 2004, ApJ, 606, 78
- Dai, X., Chartas, G., Agol, E., Bautz, M. W., Garmire, G. P. 2003, ApJ, 589, 100
- Jovanović, P., Popović, L. Č., Dimitrijević, M. S. 2003, POBeo, 76, 205
- Jovanović, P. 2005, Mem. Soc. Astron. Italiana, 7, 56
- Jovanović, P., Popović, L. Č. 2005, PASRB, 5, 195
- Jovanović, P., Popović, L. Č., Zakharov, F. 2006, in preparation
- Popović, L. Č., Mediavilla, E. G., Muñoz, J. A., Dimitrijević, M. S., Jovanović, P. 2001, SerAJ, 164, 53
- Popović, L. Č., Jovanović, P. 2002, POBeo, 73, 215
- Popović, L. Č., Jovanović, P., Bon, E., Dimitrijević, M. S. 2002a, POBeo, 73, 49
- Popović, L. Č., Mediavilla, E. G., Kubičela, A. , Jovanović, P. 2002b, A&A, 390, 473
- Popović, L. Č., Jovanović, P., Mediavilla, E. G., Muñoz, J. A. 2003a, A&AT, 22, 719
- Popović, L. Č., Mediavilla, E. G., Jovanović, P., Muñoz, J. A. 2003b, A&A, 398, 975
- Popović, L. Č., Jovanović, P., Mediavilla, E. G., Zakharov, A. F., Abajas, C., Muñoz, J. A., Chartas, G. 2006, ApJ, 637, 620
- Zakharov, F., Popović, L. Č., Jovanović, P. 2004, A&A, 420, 881
- Zakharov, F., Popović, L. Č., Jovanović, P. 2005, IAUS, 225, 363




Fig. 1.— Illustration of the straight-fold caustic crossing over an accretion disk (left), the corresponding deformations of the X-ray continuum (middle) and the Fe $K\alpha$ line (right). Undeformed profiles are presented by solid and deformed by dotted lines.




Fig. 2.— The variations of normalized total continuum flux in optical (3500-7000 Å), UV (1000-3500 Å) and X (1.24-12.4 Å i.e. 1-10 KeV) band due to microlensing by a caustic crossing over accretion disk in Schwarzschild metric. Time scale corresponds to the QSO 2237+0305 gravitational lens, where redshifts of microlens and source are $z_l = 0.04$ and $z_d = 1.69$, respectively (Jovanović et al. 2006).